



Achilles Tendon Rupture Nonoperative Rehab Protocol Prescription

Patient Name: _____

Date: _____

Diagnosis: *Achilles tendon rupture L / R ankle*

Number of visits each week: 1 2 3 4

Treatment duration _____ weeks

0-2 weeks:

Splint in plantar flexion

Non-weightbearing with crutches

2-4 weeks:

Walking boot with 20° plantar flexion heel lift* †

Non-weightbearing with crutches

Remove boot in seated position with lower extremity hanging free off table/chair/bed

Active ankle dorsiflexion to neutral only followed by passive gravity-assisted plantar flexion as far as is comfortable only

Modalities as indicated

Daily HEP for active dorsiflexion and passive plantar flexion as detailed above

5 minutes every hour

4-6 weeks:

Remove heel lift from boot

Non-weightbearing in boot

Continue 2-4 week exercise protocol

6-8 weeks:

Weightbearing in boot with crutches

May remove boot for sleeping

Continue same exercise protocol

8-12 weeks:

Wean off boot first then crutches

Initiate toe-raising exercises using the unaffected leg to support the injured leg

Once able to perform toe-raises with the injured leg unsupported, may begin Achilles stretching, strengthening and proprioception exercises



Hospital for Joint Diseases

NYU LANGONE MEDICAL CENTER

Laith M. Jazrawi, M.D.

Chief, Division of Sports Medicine

Associate Professor of Orthopaedic Surgery

Tel: (646) 501-7223 option 4, option 2

Fax: (646) 501-7234

Web: newyorkortho.com

orthosurgery.med.nyu.edu/sports-medicine

>12 weeks:

Continue to progress range of motion, strength, proprioception

Retrain strength, power, endurance

Increase dynamic weight-bearing exercise, including plyometric training

Sport-specific retraining

* Patients required to wear the boot while sleeping for first 6 weeks.

† Patients can remove the boot for bathing and dressing, but are required to adhere to the weightbearing restrictions according to the rehabilitation protocol.

Comments:

Signature: _____

Date: _____