



Associate Professor of Orthopaedics Chief - Division of Sports Medicine Tel: (212) 598-6784

Rehabilitation Protocol: Arthroscopic Meniscectomy/Chondral Debridement

Name:	Date:
Diagnosis:	Date of Surgery:
Diagnosis:	Date of Surgery.
Phase I (Weeks 0-2)	
• Weightbearing: As tolerated with crutches (for balance	e) x 24-48 hours – progress to WBAT
 Range of Motion – AAROM → AROM as tolerated 	
 Goal: Immediate full range of motion 	
 Therapeutic Exercises 	
 Quad and Hamstring sets 	
 Heel slides 	
 Co-contractions 	
 Isometric adduction and abduction exercises 	
 Straight-leg raises 	
 Patellar mobilization 	
Phase II (Weeks 2-4)	
• Weightbearing: As tolerated	
• Range of Motion – maintain full ROM – gentle passive s	stretching at end ranges
Therapeutic Exercises	vv cooming at one ranges
Quadriceps and Hamstring strengthening	
o Lunges	
Wall-sits	
 Balance exercises – Core work 	
Phase III (Weeks 4-6)	
Weightbearing: Full weightbearing	
 Range of Motion – Full/Painless ROM 	
 Therapeutic Exercises 	
 Leg press 	
 Hamstring curls 	
 Squats 	
 Plyometric exercises 	
 Endurance work 	
 Return to athletic activity as tolerated 	
Comments:	
Frequency: times per week	weeks
Signature:	Date: