

Associate Professor of Orthopaedics Chief - Division of Sports Medicine Tel: (212) 598-6784

Rehabilitation Protocol:

ACL and MCL Reconstruction				
Name:	Date:			
Diagnosis:	Date of Surgery:			
EARLY PH	ASE (Weeks 0-4)			
• Weigl	nt Bearing and Range of Motion			
0	0-6 weeks: toe-touch weight bearing w/ crutches			
0	ROM: A/AAROM 0-90° as tolerated			
 Brace 	Use:			
0	Locked in full extension at all times other than PT			
Thera	peutic Elements:			
0	Modalities as needed			
0	Patella Mob; SLR's with electric stim.; co-contractions, prone hangs			
0	 Estim; Cocontractions 			
0	No abduction of hip or leg at any time.			
0	No prone hangs if PCL reconstruction!!			
 Goals 				
	a/aa/ROM: 0-0-90			
0	Control pain/swelling			
0	Quad control			
RECOVER	RY PHASE (Weeks 5-8)			
 Weight 	nt Bearing and Range of Motion:			
0	Discontinue crutches at week 6			
 Brace 	Use:			
	At all times, open to AROM; discontinue at week 8			
 Thera 	peutic Elements:			
0	Continue above			
0	Gentle hip abduction with no resistance below knee			

- - o Wall-sits 0-45
 - o Mini-squats with support 0-45
 - Carpet drags (not with PCL reconstruction!!)
 - o Pool therapy
 - o Treadmill walking by 8 weeks
- Goals:
 - a/aa/ROM: 0-0-110 by 6 weeks and free by 8 weeks
 - o SLR x 30
 - No effusion

STRENGTHEN PHASE (Weeks 8-12)

- Weight Bearing and Range of Motion:
 - o Full
- **Therapeutic Elements:**
 - o Continue above with increased resistance
 - o Step-downs
 - Treadmill
 - $\circ \quad Stretching \\$

Laith M. Jazrawi, MD



Associate Professor of Orthopaedics Chief - Division of Sports Medicine Tel: (212) 598-6784

0	Begin prone hangs and HSL (if PCL reconstruction)
Goals:	

o Walk 1-2 miles at 15 min/mile pace

	REINTEGRATION PHASE (Months 3-5)
-	 Weight Bearing and Range of Motion:

- o Full
- Brace Use:
 - o None
 - o If return to sport, fitting for custom brace by 5 months
 - o Can start jogging/running at 6 months
- Therapeutic Elements:
 - o Slide boards
 - o Begin agility drills
 - o Figure 8's
 - o Gentle loops
 - o Large zig-zags
 - \circ Swimming
 - o Begin plyometrics at 4 months
- Goals:
 - o Treadmill (walk 1-2 miles at 10-12 min/mile pace)
 - o Return to competitive activities

•				
Co	m	m	Δn	TC:
w				L.7.

Frequency: times per week	Duration: weeks
Signature:	Date: