

## PRP Therapy Instructions

### Before Arriving for Your Exam

You will undergo complete blood count (CBC) testing to determine if your platelet count is sufficient. Do not take aspirin or aspirin-related medications such as non-steroidal anti-inflammatory drugs (NSAID), such as Aleve® or Advil®, for **1 week before** your therapy and for **2 weeks after**. You should discuss any changes to your medications with your physician.

**These include:**

- Abciximab (ReoPro) NSAIDs (ibuprofen, naproxen, aspirin)
- Cilostazol (Pletal) Prasugrel (Effient)
- Clopidogrel (Plavix) Ticagrelor (Brilinta)
- Dipyridamole (Persantine) Ticlopidine (Ticlid)
- Eptifibatide (Integrilin) Tirofiban (Aggrastat)

**Stop taking supplements affecting platelets for two weeks before PRP therapy.**

**These include:**

- Black currant seed oil Krill oil
- Borage oil Nattokinase
- Bromelain Papain
- Evening primrose oil Serrapeptase
- Fish oil Wobenzyme
- Flaxseed oil

You should not have PRP therapy if you have:

- Abnormal platelet function
- Active systemic infection
- Active cancer
- Low-platelet count
- Severe anemia

### After Arriving

A technologist will ask you a few questions regarding your medical history. They will also do a few simple tests such as taking your blood pressure.

Please inform the technologist and/or physician of any allergies you may have before your exam. A physician will discuss this therapy with you in detail and answer any questions you may have.

Immediately before your scheduled PRP therapy, blood is collected and the platelets, which contain your body's natural healing components (growth factors), are concentrated in a centrifuge.



## During Your Exam

During the treatment, a local anesthetic is used on the affected region and then the platelet rich plasma is reinjected under ultrasound-guidance to stimulate and enhance healing.

In most cases one treatment is enough, but in certain cases between one and three treatments are recommended for maximum benefit over 4 to 6 weeks.

## After Your Exam

### PRP Post-Operative Instructions:

Immediately after the procedure, you will remain lying down and under observation for a few minutes up to 30 minutes.

Do not take aspirin or any aspirin-related drugs **for two weeks after your injection**. Your Musculoskeletal Imaging Physician will prescribe pain medication as needed.

1. Apply an ice pack for 10-15 minutes on the injected area every 2 hours for 6-8 hours following the procedure.
2. You may apply a lidoderm patch for 6-7 hours for up to 12 hours after the procedure.
3. Avoid major physical activity for two to four weeks.
4. Do not take any drugs or food supplements (see list above) affecting platelets for one to two weeks.

Avoid heavy repetitive activities to affected area for 4 weeks.

Patients typically notice improvement in 6 weeks

If the affected area is your foot, ankle, or knee, you will need to wear a boot-type immobilizer. For 48 hours you will need to keep all weight off that foot. The boot immobilizer should be worn for 2 to 3 weeks after the injection.

Avoid heavy or repetitive physical activity with the treated body part for up to a week to allow the healing to occur. If the affected area is your elbow, for example, avoid vigorous activities such as strong gripping or lifting of loads heavier than 10 pounds for 4 weeks after the injection.

There may be some swelling and soreness at the injection site for a day or two. Ice and pain medicine should alleviate the soreness and swelling. Elevate the leg or arm and limit your activities as much as needed to remain comfortable.

You might have some discomfort in the area of the injection that can last a few days or up to a week. Sometimes it can seem like your injury is worse than before the treatment, but that is because an inflammatory response has just been stimulated. This is common, and the temporary worsening of your symptoms usually doesn't last. Since PRP's effectiveness is based on your own body's ability to heal, most patients notice improvement after 2 to 6 weeks. In most cases, pain is replaced with soreness and then the soreness goes away. Strength and endurance slowly increase with time. Many patients notice improvement for as long as 6 to 9 months.