



Associate Professor of Orthopaedics Chief - Division of Sports Medicine Tel: (212) 598-6784

Rehabilitation Protocol: Meniscus Allograft Transplantation

Name:	Date:
Diagnosis: _	Date of Surgery:
Phase I (W	Veeks 0-8)
-	htbearing:
_	Weeks 0-2: Partial Weightbearing (up to 50%)
	Weeks 2-6: Advance to WBAT with crutches (d/c crutches at 4 weeks post-op if gait normalized)
	ed Knee Brace: worn for 6 weeks post-op
_	Locked in full extension for ambulation and sleeping – remove for hygiene (Week 1)
0	
0	
0	
	e of Motion – PROM → AAROM → AROM as tolerated
U	Weeks 0-2 : Non-weightbearing 0-90°
	Weeks 2-8: Full non-weightbearing ROM as tolerated – progress to flexion angles greater than 90°
	apeutic Exercises
	Quadriceps sets, heel slides, straight leg raises, patellar mobilizations, co-contractions (Weeks 0-2)
0	Add heel raises and terminal knee extensions (Weeks 2-8) Activities in brace for first 6 weeks – then without brace
0	
0	No weightbearing with flexion > 90° during Phase I
0	Avoid tibial rotation for first 8 weeks to protect the meniscal allograft
Phase II	(Weeks 8-12)
	htbearing: As tolerated
	e of Motion – Full active ROM
_	apeutic Exercises
	Progress to closed chain extension exercises, begin hamstring strengthening
	Lunges – 0-90°, Leg press – 0-90° (flexion only)
0	
O	begin use of the stationary bicycle
Phase III	(Months 3-6)
• Weig	htbearing: Full weightbearing with normal gait pattern
_	e of Motion - Full/Painless ROM
U	apeutic Exercises
	Continue with quad and hamstring strengthening
	Begin jogging/running
	Plyometrics and sport-specific drills
	ual return to athletic activity as tolerated (6 months post-op)
	tenance program for strength and endurance
Manie	chance program for strength and chaurance
Comments:	
Frequency:	times per week Duration: weeks
Signature:	Date