

Proximal or Distal Realignment +/- Medial Patellofemoral Ligament Reconstruction Rehabilitation Program

Name	:		Date:
Diagn	osis:		Date of Surgery:
Pha	se I –In	nmediate Postoperative Phase (Day 1-5)	
•	Goals:		
	0	Diminish swelling/inflammation (control hemarth	rosis)
	0	Diminish postoperative pain	•
	0	Initiate voluntary quadriceps control	
	0	Independent ambulation	
•	Brace:		
	0	Brace for ambulation only (POD 1 to Week 4)	
•	Weigh	t-bearing	
	0	As tolerated with two crutches (approx. 50% WB)	
•	Swelli	ng/Inflammation Control	
	0	Cryotherapy	
	0	Compression bandages	
	0	Elevation & ankle pumps	
•	Range	of Motion	
	0	Full passive knee extension	
	0	Flexion to 45° (day 1-4)	
	0	Flexion to 60° (day 5)	
	0	PROM and gentle AAROM only	
•	Flexibi	ility	
	0	Hamstring and calf stretches	
	0	PROM/AAROM within ROM limitations	
Ph	ase II -	Acute Phase (Week 2-4)	
	Goals:		
	0	Control swelling and pain	
	0	Promote healing of realignment tibial tuberosity	
	0	Quadriceps strengthening	
•	Brace		
	0	Continue brace for ambulation only	
	0	Discontinue brace (week 4)	
•	Weigh	t-bearing	
	0	Progress WBAT (2 crutches for 4 weeks)	
•	Swelli	ng/inflammation	
	0	Continue use of cryotherapy	

Range of motion

o Elevation

- o PROM/AAROM exercises
- o ROM 0-75° (week 1-3)

o Compression bandage

- o ROM 0-90° (week 4)
- Muscle Retraining
 - o Electrical muscle stimulation to quads

Associate Professor of Orthopaedics Chief - Division of Sports Medicine Tel: (212) 598-6784

- Quad setting isometrics
- Straight leg raises (flexion)
- Hip adduction/abduction
- Hip extension
- o GENTLE submaximal isometric knee extension
- o Week 4
 - Light leg press
 - Vertical squats (no weight)
- Flexibility
 - o Continue hamstring, calf stretches

Phase III -Subacute Phase "Motion" Phase (Week 5-8)

- Goals
 - Gradual improvement in ROM
 - o Improve muscular strength and endurance
 - o Control forces on extension mechanism
- Weight-bearing
 - o One crutch (week 4-6)
 - o Discontinue crutch (week 6)
- Range of motion
 - o PROM 0-115°
 - o PROM 0-125°
 - o PROM 0-125/135°
- Exercises
 - Continue electrical muscle stimulation to quadriceps
 - Quadriceps setting isometric
 - o Hip adduction, abuction, and extension
 - Vertical squats
 - Leg press
 - o Knee extension light (0-60°)
 - o Bicycle (week 6-8)
 - Pool program [walking, strengthening (when able)]
- Flexibility
 - Continue all stretching exercises for LE

Phase IV -Strengthening Phase (weeks 9-16)

- Criteria to Progress to Phase IV
 - o ROM at least 0-115 degrees
 - Absence of swelling/inflammation
 - Voluntary control of quads
- Goals
 - o Gradual improvement of muscular strength
 - Functional activities/drills
- Exercises
 - o ½ vertical squats (0-60°)
 - o Wall squats (0-60°)
 - Leg press
 - o Forward lunges
 - o Lateral lunges
 - o Lateral step-ups
 - Knee extension (60-0°)



Laith M. Jazrawi, MD Associate Professor of Orthopaedics

Chief - Division of Sports Medicine
Tel: (212) 598-6784

- o Hip adduction/abduction
- o Bicycle
- o Stairmaster

Phase V -Return to Activity Pha

- Criteria to enter Phase V
 - o Full non-painful ROM
 - Appropriate strength level (80% or greater of contralateral leg)
 - o Satisfactory clinical exam
- Goals

Comments:

- o Functional return to specific drills
- Exercises
 - o Functional drills
 - Strengthening exercises
 - o Flexibility exercises

Frequency: times per week	Duration: weeks
	_