

Post-Operative Instructions

Tibial Plateau Fracture Arthroscopic Reduction and Fixation

Day of surgery

- A. Diet as tolerated
- B. Icing is important for the first 5-7 days post-op. While the post-op dressing is in place, icing should be done continuously. Once the dressing is removed on the third post-operative day, ice is applied for 20-minute periods 3-4 times per day. Care must be taken with icing to avoid frostbite. Alternatively, Cryocuff or Game-ready ice cuff can be used as per instructions.
- C. Pain medication as needed every 4-6 hours (refer to pain medication sheet).
- D. Make sure you have a physical therapy post-op appointment scheduled during the first week after surgery.

Video instructions for your brace can be found at <https://www.youtube.com/watch?v=jyRZkSyFBOQ>

First Post-Operative Day

- A. Continue icing
- B. Pain medication as needed.
- C. If you have been assigned a Continuous Passive Motion (CPM) machine, it should be started during the first week after your surgery. This machine will be set at 30 degrees. The machine should be used 6 hours per day (2 hours in the morning, 2 hours in the afternoon, and 2 hours in the evening). You will use this machine for 1 month after surgery. Do not wear leg brace or cooling device while using CPM machine.

Second Post-Operative Day Until Return Visit

- A. Continue icing
- B. Unless otherwise noted, no weightbearing for the first 6 weeks after surgery. After 6 weeks, you can bear as much weight on the affected leg as you can tolerate.
- C. Call our office @ 646-501-7223 option 4, option 2 to confirm your first postoperative visit, which is usually about 1-2 weeks after surgery if you have not been given a time. If you are experiencing any problems, please call our office or contact us via the internet at www.newyorkortho.com.

Third Post-Operative Day

- A. You may remove surgical bandage and shower this evening. Apply 4x4 (or similar size) waterproof bandage to these wounds prior to showering and when showering is complete apply fresh waterproof bandage. Please ensure that the bandage is large enough to completely cover the incision. You will need to follow this routine for 2 weeks after surgery.

Rehabilitation Protocol: Tibial Plateau Fracture Arthroscopic Reduction and Fixation

Name: _____ Date: _____

Diagnosis: _____ Date of Surgery: _____

Phase I (Weeks 0-4)

- **Weightbearing:** Toe touch weight bearing x 4 weeks
- **Hinged Knee Brace:**
 - Locked in full extension for ambulation and sleeping (**Weeks 0-4**)
- **Range of Motion** – AAROM → AROM as tolerated
- **Therapeutic Exercises**
 - Quad/Hamstring sets
 - Heel slides
 - Non-weightbearing stretch of the Gastroc/Soleus
 - Straight-Leg Raise with brace in full extension until quad strength prevents extension lag

Phase II (Weeks 4-8)

- **Weightbearing:** Start partial weightbearing and progress to full weightbearing by 6 weeks without crutches
- **Hinged Knee Brace:**
 - Unlock brace Week 6
 - Discontinue brace use when patient has achieved full extension with no evidence of extension lag by week 8
- **Range of Motion** – Maintain full knee extension – work on progressive knee flexion
- **Therapeutic Exercises**
 - Closed chain extension exercises
 - Hamstring curls
 - Toe raises
 - Balance exercises
 - Progress to weightbearing stretch of the Gastroc/Soleus
 - Begin use of the stationary bicycle

Phase III (Weeks 8-16)

- **Weightbearing:** Full weightbearing
- **Range of Motion** – Full/Painless ROM
- **Therapeutic Exercises**
 - Advance closed chain strengthening exercises, proprioception activities
 - Begin use of the Stairmaster/Elliptical
 - **Can Start Straight Ahead Running at 12 Weeks**

Phase IV (Months 4-6)

- Gradual return to athletic activity as tolerated
- Maintenance program for strength and endurance

Comments:

Frequency: ____ times per week

Duration: ____ weeks

Signature: _____

Date: _____